

Package ‘Copula.surv’

May 15, 2024

Type Package

Title Analysis of Bivariate Survival Data Based on Copulas

Version 1.5

Date 2024-05-15

Author Takeshi Emura

Maintainer Takeshi Emura <takeshiemura@gmail.com>

Description Simulating bivariate survival data from copula models.

Estimation of the association parameter in copula models.

Two different ways to estimate the association parameter in copula models are implemented.

A goodness-of-fit test for a given copula model is implemented.

See Emura, Lin and Wang (2010) <[doi:10.1016/j.csda.2010.03.013](https://doi.org/10.1016/j.csda.2010.03.013)> for details.

License GPL-2

NeedsCompilation no

Repository CRAN

Date/Publication 2024-05-15 04:50:03 UTC

R topics documented:

Copula.surv-package	2
simu.BB1	2
simu.CC	4
simu.Clayton	5
simu.FGM	6
simu.Frank	7
simu.GB	8
simu.Gumbel	10
Test.Clayton	11
Test.Gumbel	12
U1.Clayton	13
U1.Gumbel	15
U2.Clayton	16
U2.Gumbel	17

Weib.reg.BB1	18
Weib.reg.BB1.0	19
Weib.reg.Clayton	20

Index**22**

Copula.surv-package *Analysis of Bivariate Survival Data*

Description

Simulating bivariate survival data from copula models (Emura et al. 2019). Estimation of the association parameter in copula models. Two different ways to estimate the association parameter in copula models are implemented. A goodness-of-fit test for a given copula model is implemented. See Emura, Lin and Wang (2010) <doi:10.1016/j.csda.2010.03.013> for details. Also, Weibull regression is implemented (Section 2.6.3 of Emura et al. (2019)).

Details

Details are seen from the references.

Author(s)

Takeshi Emura Maintainer: Takeshi Emura <takeshiemura@gmail.com>

References

- Emura T, Lin CW, Wang W (2010) A goodness-of-fit test for Archimedean copula models in the presence of right censoring, Compt Stat Data Anal 54: 3033-43
 Emura T, Matsui S, Rondeau V (2019), Survival Analysis with Correlated Endpoints, Joint Frailty-Copula Models, JSS Research Series in Statistics, Springer

simu.BB1

Simulating data from the BB1 copula

Description

n pairs of (U,V) are generated from the BB1 copula. n pairs of (X,Y) are generated from the corresponding bivariate survival model with the Weibull marginal distributions. The default parameters (scale1=scale2=shape1=shape2=1) give the unit exponential distributions.

Usage

```
simu.BB1(n,alpha,d=0,scale1=1,scale2=1,shape1=1,shape2=1,Print=FALSE)
```

Arguments

n	sample size
alpha	association (copula) parameter
d	BB1 copula's departure parameter from the Clayton (d=0 is the default)
scale1	scale parameter for X
scale2	scale parameter for Y
shape1	shape parameter for X
shape2	shape parameter for Y
Print	print Kendall's tau and means of X and Y if "TRUE"

Details

See Section 2.6 of Emura et al.(2019) for copulas and bivariate survival times.

Value

U	uniformly distributed on (0,1)
V	uniformly distributed on (0,1)
X	Weibull distributed (scale1, shape1)
Y	Weibull distributed (scale2, shape2)

Author(s)

Takeshi Emura

References

Emura T, Lin CW, Wang W (2010) A goodness-of-fit test for Archimedean copula models in the presence of right censoring, Compt Stat Data Anal 54: 3033-43

Emura T, Matsui S, Rondeau V (2019), Survival Analysis with Correlated Endpoints, Joint Frailty-Copula Models, JSS Research Series in Statistics, Springer

Examples

```

n=100
Dat=simu.BB1(n=n,alpha=1,d=2,scale1=1,scale2=2,shape1=0.5,shape2=2)
plot(Dat[, "U"],Dat[, "V"])
cor(Dat[, "U"],Dat[, "V"],method="kendall")
plot(Dat[, "X"],Dat[, "Y"])
cor(Dat[, "X"],Dat[, "Y"],method="kendall")

```

simu.CC*Simulating data from the Celebioglu-Cuadras (CC) copula***Description**

n pairs of (U,V) are generated from the CC copula. n pairs of (X,Y) are generated from the corresponding bivariate survival model with the Weibull marginal distributions. The default parameters (scale1=scale2=shape1=shape2=1) give the unit exponential distributions.

Usage

```
simu.CC(n,alpha,scale1=1,scale2=1,shape1=1,shape2=1,Print=FALSE)
```

Arguments

n	sample size
alpha	association (copula) parameter, -1<=alpha<=1
scale1	scale parameter for X
scale2	scale parameter for Y
shape1	shape parameter for X
shape2	shape parameter for Y
Print	print Kendall's tau and means of X and Y if "TRUE"

Details

See Section 2.6 of Emura et al.(2019) for copulas and bivariate survival times.

Value

U	uniformly distributed on (0,1)
V	uniformly distributed on (0,1)
X	Weibull distributed (scale1, shape1)
Y	Weibull distributed (scale2, shape2)

Author(s)

Takeshi Emura

References

- Emura T, Lin CW, Wang W (2010) A goodness-of-fit test for Archimedean copula models in the presence of right censoring, Compt Stat Data Anal 54: 3033-43
- Emura T, Matsui S, Rondeau V (2019), Survival Analysis with Correlated Endpoints, Joint Frailty-Copula Models, JSS Research Series in Statistics, Springer

Examples

```
n=100
Dat=simu.CC(n=n,alpha=-1,scale1=1,scale2=2,shape1=0.5,shape2=2)
plot(Dat[,"U"],Dat[,"V"])
cor(Dat[,"U"],Dat[,"V"],method="kendall")
plot(Dat[,"X"],Dat[,"Y"])
cor(Dat[,"X"],Dat[,"Y"],method="kendall")
```

simu.Clayton

Simulating data from the Clayton copula

Description

n pairs of (U, V) are generated from the Clayton copula. n pairs of (X, Y) are generated from the corresponding bivariate survival model with the Weibull marginal distributions. The default parameters ($scale1=scale2=shape1=shape2=1$) give the unit exponential distributions.

Usage

```
simu.Clayton(n,alpha,scale1=1,scale2=1,shape1=1,shape2=1,Print=FALSE)
```

Arguments

n	sample size
α	association (copula) parameter
$scale1$	scale parameter for X
$scale2$	scale parameter for Y
$shape1$	shape parameter for X
$shape2$	shape parameter for Y
$Print$	print Kendall's tau and means of X and Y if "TRUE"

Details

See Section 2.6 of Emura et al.(2019) for copulas and bivariate survival times.

Value

U	uniformly distributed on $(0,1)$
V	uniformly distributed on $(0,1)$
X	Weibull distributed ($scale1$, $shape1$)
Y	Weibull distributed ($scale2$, $shape2$)

Author(s)

Takeshi Emura

References

- Emura T, Lin CW, Wang W (2010) A goodness-of-fit test for Archimedean copula models in the presence of right censoring, *Compt Stat Data Anal* 54: 3033-43
- Emura T, Matsui S, Rondeau V (2019), Survival Analysis with Correlated Endpoints, Joint Frailty-Copula Models, JSS Research Series in Statistics, Springer

Examples

```
n=100
Dat=simu.Clayton(n=n,alpha=1,scale1=1,scale2=2,shape1=0.5,shape2=2)
plot(Dat[, "U"],Dat[, "V"])
cor(Dat[, "U"],Dat[, "V"],method="kendall")
plot(Dat[, "X"],Dat[, "Y"])
cor(Dat[, "X"],Dat[, "Y"],method="kendall")
```

simu.FGM

Simulating data from the FGM copula

Description

n pairs of (U,V) are generated from the FGM copula. n pairs of (X,Y) are generated from the corresponding bivariate survival model with the Weibull marginal distributions. The default parameters (scale1=scale2=shape1=shape2=1) give the unit exponential distributions.

Usage

```
simu.FGM(n,alpha,scale1=1,scale2=1,shape1=1,shape2=1,Print=FALSE)
```

Arguments

n	sample size
alpha	association (copula) parameter; -1<=alpha<=1
scale1	scale parameter for X
scale2	scale parameter for Y
shape1	shape parameter for X
shape2	shape parameter for Y
Print	print Kendall's tau and means of X and Y if "TRUE"

Details

See Section 2.6 of Emura et al.(2019) for copulas and bivariate survival times.

Value

U	uniformly distributed on (0,1)
V	uniformly distributed on (0,1)
X	Weibull distributed (scale1, shape1)
Y	Weibull distributed (scale2, shape2)

Author(s)

Takeshi Emura

References

- Emura T, Lin CW, Wang W (2010) A goodness-of-fit test for Archimedean copula models in the presence of right censoring, *Compt Stat Data Anal* 54: 3033-43
- Emura T, Matsui S, Rondeau V (2019), Survival Analysis with Correlated Endpoints, Joint Frailty-Copula Models, JSS Research Series in Statistics, Springer

Examples

```
n=100
Dat=simu.FGM(n=n,alpha=1,scale1=1,scale2=2,shape1=0.5,shape2=2)
plot(Dat[,"U"],Dat[,"V"])
cor(Dat[,"U"],Dat[,"V"],method="kendall")
plot(Dat[,"X"],Dat[,"Y"])
cor(Dat[,"X"],Dat[,"Y"],method="kendall")
```

simu.Frank

Simulating data from the Frank copula

Description

n pairs of (U,V) are generated from the Frank copula. n pairs of (X,Y) are generated from the corresponding bivariate survival model with the Weibull marginal distributions. The default parameters (scale1=scale2=shape1=shape2=1) give the unit exponential distributions.

Usage

```
simu.Frank(n,alpha,scale1=1,scale2=1,shape1=1,shape2=1,Print=FALSE)
```

Arguments

n	sample size
alpha	association (copula) parameter
scale1	scale parameter for X
scale2	scale parameter for Y
shape1	shape parameter for X
shape2	shape parameter for Y
Print	print Kendall's tau and means of X and Y if "TRUE"

Details

See Section 2.6 of Emura et al.(2019) for copulas and bivariate survival times.

Value

U	uniformly distributed on (0,1)
V	uniformly distributed on (0,1)
X	Weibull distributed (scale1, shape1)
Y	Weibull distributed (scale2, shape2)

Author(s)

Takeshi Emura

References

- Emura T, Lin CW, Wang W (2010) A goodness-of-fit test for Archimedean copula models in the presence of right censoring, *Compt Stat Data Anal* 54: 3033-43
 Emura T, Matsui S, Rondeau V (2019), Survival Analysis with Correlated Endpoints, Joint Frailty-Copula Models, JSS Research Series in Statistics, Springer

Examples

```
n=100
Dat=simu.Frank(n=n,alpha=10,scale1=1,scale2=2,shape1=0.5,shape2=2)
plot(Dat[, "U"],Dat[, "V"])
cor(Dat[, "U"],Dat[, "V"],method="kendall")
plot(Dat[, "X"],Dat[, "Y"])
cor(Dat[, "X"],Dat[, "Y"],method="kendall")
```

Description

n pairs of (U,V) are generated from the GB copula. n pairs of (X,Y) are generated from the corresponding bivariate survival model with the Weibull marginal distributions. The default parameters (scale1=scale2=shape1=shape2=1) give the unit exponential distributions.

Usage

```
simu.GB(n,alpha,scale1=1,scale2=1,shape1=1,shape2=1,Print=FALSE)
```

Arguments

n	sample size
alpha	association (copula) parameter, $0 \leq \text{alpha} \leq 1$
scale1	scale parameter for X
scale2	scale parameter for Y
shape1	shape parameter for X
shape2	shape parameter for Y
Print	print Kendall's tau and means of X and Y if "TRUE"

Details

See Section 2.6 of Emura et al.(2019) for copulas and bivariate survival times.

Value

U	uniformly distributed on (0,1)
V	uniformly distributed on (0,1)
X	Weibull distributed (scale1, shape1)
Y	Weibull distributed (scale2, shape2)

Author(s)

Takeshi Emura

References

Emura T, Lin CW, Wang W (2010) A goodness-of-fit test for Archimedean copula models in the presence of right censoring, Compt Stat Data Anal 54: 3033-43

Emura T, Matsui S, Rondeau V (2019), Survival Analysis with Correlated Endpoints, Joint Frailty-Copula Models, JSS Research Series in Statistics, Springer

Examples

```

n=100
Dat=simu.GB(n=n,alpha=1,scale1=1,scale2=2,shape1=0.5,shape2=2)
plot(Dat[, "U"], Dat[, "V"])
cor(Dat[, "U"], Dat[, "V"], method="kendall")
plot(Dat[, "X"], Dat[, "Y"])
cor(Dat[, "X"], Dat[, "Y"], method="kendall")

```

simu.Gumbel*Simulating data from the Gumbel copula***Description**

n pairs of (U,V) are generated from the Gumbel copula. *n* pairs of (X,Y) are generated from the corresponding bivariate survival model with the Weibull marginal distributions. The default parameters (scale1=scale2=shape1=shape2=1) give the unit exponential distributions.

Usage

```
simu.Gumbel(n,alpha,scale1=1,scale2=1,shape1=1,shape2=1,Print=FALSE)
```

Arguments

<i>n</i>	sample size
<i>alpha</i>	association (copula) parameter
<i>scale1</i>	scale parameter for X
<i>scale2</i>	scale parameter for Y
<i>shape1</i>	shape parameter for X
<i>shape2</i>	shape parameter for Y
<i>Print</i>	print Kendall's tau and means of X and Y if "TRUE"

Details

See Section 2.6 of Emura et al.(2019) for copulas and bivariate survival times.

Value

<i>U</i>	uniformly distributed on (0,1)
<i>V</i>	uniformly distributed on (0,1)
<i>X</i>	Weibull distributed (scale1, shape1)
<i>Y</i>	Weibull distributed (scale2, shape2)

Author(s)

Takeshi Emura

References

- Emura T, Lin CW, Wang W (2010) A goodness-of-fit test for Archimedean copula models in the presence of right censoring, Compt Stat Data Anal 54: 3033-43
- Emura T, Matsui S, Rondeau V (2019), Survival Analysis with Correlated Endpoints, Joint Frailty-Copula Models, JSS Research Series in Statistics, Springer

Examples

```
n=100
Dat=simu.Gumbel(n=n,alpha=1,scale1=1,scale2=2,shape1=0.5,shape2=2)
plot(Dat[,"U"],Dat[,"V"])
cor(Dat[,"U"],Dat[,"V"],method="kendall")
plot(Dat[,"X"],Dat[,"Y"])
cor(Dat[,"X"],Dat[,"Y"],method="kendall")
```

Test.Clayton

A goodness-of-fit test for the Clayton copula

Description

Perform a goodness-of-fit test for the Clayton copula based on Emura, Lin and Wang (2010). The test is asymptotically equivalent to the test of Shih (1998).

Usage

```
Test.Clayton(x.obs,y.obs,dx,dy,lower=0.001,upper=50,U.plot=TRUE)
```

Arguments

x.obs	censored times for X
y.obs	censored times for Y
dx	censoring indicators for X
dy	censoring indicators for Y
lower	lower bound for the association parameter
upper	upper bound for the association parameter
U.plot	if TRUE, draw the plot of U_1(theta)

Details

See the references.

Value

theta1	association parameter by the pseudo-likelihood estimator
theta2	association parameter by the unweighted estimator
Stat	$\log(\theta_1) - \log(\theta_2)$
Z	Z-value of the goodness-of-fit for the Clayton copula
P	P-value of the goodness-of-fit for the Clayton copula

Author(s)

Takeshi Emura

References

- Emura T, Lin CW, Wang W (2010) A goodness-of-fit test for Archimedean copula models in the presence of right censoring, *Compt Stat Data Anal* 54: 3033-43
 Shih JH (1998) A goodness-of-fit test for association in a bivariate survival model. *Biometrika* 85: 189-200

Examples

```
n=20
theta_true=2 ## association parameter ##
r1_true=2 ## hazard for X
r2_true=2 ## hazard for Y

set.seed(1)
V1=runif(n)
V2=runif(n)
X=-1/r1_true*log(1-V1)
W=(1-V1)^(-theta_true)
Y=1/theta_true/r2_true*log( 1-W+W*(1-V2)^(-theta_true/(theta_true+1)) )
C=runif(n,min=0,max=5)

x.obs=pmin(X,C)
y.obs=pmin(Y,C)
dx=X<=C
dy=Y<=C

Test.Clayton(x.obs,y.obs,dx,dy)
```

Test.Gumbel

A goodness-of-fit test for the Gumbel copula

Description

Perform a goodness-of-fit test for the Gumbel copula based on Emura, Lin and Wang (2010).

Usage

```
Test.Gumbel(x.obs,y.obs,dx,dy,lower=0.01,upper=50,U.plot=TRUE)
```

Arguments

x.obs	censored times for X
y.obs	censored times for Y
dx	censoring indicators for X
dy	censoring indicators for Y
lower	lower bound for the association parameter
upper	upper bound for the association parameter
U.plot	if TRUE, draw the plot of U_1(theta) and U_2(theta)

Details

See the references.

Value

theta1	association parameter by the pseudo-likelihood estimator
theta2	association parameter by the unweighted estimator
Stat	$\log(\text{theta1}) - \log(\text{theta2})$
Z	Z-value of the goodness-of-fit for the Clayton copula
P	P-value of the goodness-of-fit for the Clayton copula

Author(s)

Takeshi Emura

References

Emura T, Lin CW, Wang W (2010) A goodness-of-fit test for Archimedean copula models in the presence of right censoring, *Compt Stat Data Anal* 54: 3033-43

Examples

```
x.obs=c(1,2,3,4,5,6,7,8,9,10,11,12,13,14,15)
y.obs=c(2,1,4,5,6,8,3,7,10,9,11,12,13,14,15)
dx=c(1,1,1,1,1,1,1,1,1,1,1,1,1,1,1)
dy=c(1,1,1,1,1,1,1,1,1,1,1,1,1,1,1)
Test.Gumbel(x.obs,y.obs,dx,dy)
```

Description

Estimate the association parameter of the Clayton copula using bivariate survival data. The estimator was derived by Clayton (1978) and reformulated by Emura, Lin and Wang (2010).

Usage

```
U1.Clayton(x.obs,y.obs,dx,dy,lower=0.001,upper=50,U.plot=TRUE)
```

Arguments

x.obs	censored times for X
y.obs	censored times for Y
dx	censoring indicators for X
dy	censoring indicators for Y
lower	lower bound for the association parameter
upper	upper bound for the association parameter
U.plot	if TRUE, draw the plot of U_1(theta)

Details

Details are seen from the references.

Value

theta	association parameter
tau	Kendall's tau (=theta/(theta+2))

Author(s)

Takeshi Emura

References

Clayton DG (1978). A model for association in bivariate life tables and its application to epidemiological studies of familial tendency in chronic disease incidence. Biometrika 65: 141-51.

Emura T, Lin CW, Wang W (2010) A goodness-of-fit test for Archimedean copula models in the presence of right censoring, Compt Stat Data Anal 54: 3033-43

Examples

```

n=200
theta_true=2 ## association parameter ##
r1_true=1 ## hazard for X
r2_true=1 ## hazard for Y

set.seed(1)
V1=runif(n)
V2=runif(n)
X=-1/r1_true*log(1-V1)
W=(1-V1)^(-theta_true)
Y=1/theta_true/r2_true*log( 1-W+W*(1-V2)^(-theta_true/(theta_true+1)) )
C=runif(n,min=0,max=5)

x.obs=pmin(X,C)
y.obs=pmin(Y,C)
dx=X<=C
dy=Y<=C

```

```
U1.Clayton(x.obs,y.obs,dx,dy)
```

U1.Gumbel

Estimation of an association parameter via the unweighted estimator

Description

Estimate the association parameter of the Gumbel copula using bivariate survival data. The estimator was derived by Emura, Lin and Wang (2010).

Usage

```
U1.Gumbel(x.obs,y.obs,dx,dy,lower=0.01,upper=50,U.plot=TRUE)
```

Arguments

x.obs	censored times for X
y.obs	censored times for Y
dx	censoring indicators for X
dy	censoring indicators for Y
lower	lower bound for the association parameter
upper	upper bound for the association parameter
U.plot	if TRUE, draw the plot of U_1(theta)

Details

Details are seen from the references.

Value

theta	association parameter
tau	Kendall's tau (=theta/(theta+2))

Author(s)

Takeshi Emura

References

Emura T, Lin CW, Wang W (2010) A goodness-of-fit test for Archimedean copula models in the presence of right censoring, Compt Stat Data Anal 54: 3033-43

Examples

```
x.obs=c(1,2,3,4,5,6,7,8,9,10,11,12,13,14,15)
y.obs=c(2,1,4,5,6,8,3,7,10,9,11,12,13,14,15)
dx=c(1,1,1,1,1,1,1,1,1,1,1,1,1,1,1)
dy=c(1,1,1,1,1,1,1,1,1,1,1,1,1,1,1)
U1.Gumbel(x.obs,y.obs,dx,dy)
```

U2.Clayton

Estimation of an association parameter via the unweighted estimator

Description

Estimate the association parameter of the Clayton copula using bivariate survival data. The estimator was defined as the unweighted estimator in Emura, Lin and Wang (2010).

Usage

```
U2.Clayton(x.obs,y.obs,dx,dy)
```

Arguments

x.obs	censored times for X
y.obs	censored times for Y
dx	censoring indicators for X
dy	censoring indicators for Y

Details

Details are seen from the references.

Value

theta	association parameter
tau	Kendall's tau (=theta/(theta+2))

Author(s)

Takeshi Emura

References

Emura T, Lin CW, Wang W (2010) A goodness-of-fit test for Archimedean copula models in the presence of right censoring, Compt Stat Data Anal 54: 3033-43

Examples

```

n=200
theta_true=2 ## association parameter ##
r1_true=1 ## hazard for X
r2_true=1 ## hazard for Y

set.seed(1)
V1=runif(n)
V2=runif(n)
X=-1/r1_true*log(1-V1)
W=(1-V1)^(-theta_true)
Y=1/theta_true/r2_true*log( 1-W+W*(1-V2)^(-theta_true/(theta_true+1)) )
C=runif(n,min=0,max=5)

x.obs=pmin(X,C)
y.obs=pmin(Y,C)
dx=X<=C
dy=Y<=C

U2.Clayton(x.obs,y.obs,dx,dy)

```

U2.Gumbel

Estimation of an association parameter via the pseudo-likelihood

Description

Estimate the association parameter of the Gumbel copula using bivariate survival data. The estimator was derived by Emura, Lin and Wang (2010).

Usage

```
U2.Gumbel(x.obs,y.obs,dx,dy,lower=0.01,upper=50,U.plot=TRUE)
```

Arguments

x.obs	censored times for X
y.obs	censored times for Y
dx	censoring indicators for X
dy	censoring indicators for Y
lower	lower bound for the association parameter
upper	upper bound for the association parameter
U.plot	if TRUE, draw the plot of U_1(theta)

Details

Details are seen from the references.

Value

theta	association parameter
tau	Kendall's tau (=theta/(theta+1))

Author(s)

Takeshi Emura

References

Emura T, Lin CW, Wang W (2010) A goodness-of-fit test for Archimedean copula models in the presence of right censoring, Compt Stat Data Anal 54: 3033-43

Examples

```
x.obs=c(1,2,3,4,5)
y.obs=c(2,1,4,5,6)
dx=c(1,1,1,1,1)
dy=c(1,1,1,1,1)
U2.Gumbel(x.obs,y.obs,dx,dy)
```

Weib.reg.BB1

Weibull regression under the BB1 copula

Description

See Section 2.6.3 of Emura et al. (2019).

Usage

```
Weib.reg.BB1(x.obs,y.obs,dx,dy,zx,zy,convergence.par=FALSE)
```

Arguments

x.obs	censored times for X
y.obs	censored times for Y
dx	censoring indicators for X
dy	censoring indicators for Y
zx	matrix of covariates for X
zy	matrix of covariates for Y
convergence.par	if TRUE, show the details

Details

Details are seen from the references.

Value

beta_x	regression coefficients for X
beta_y	regression coefficients for Y

Author(s)

Takeshi Emura

References

Emura T, Matsui S, Rondeau V (2019), Survival Analysis with Correlated Endpoints, Joint Frailty-Copula Models, JSS Research Series in Statistics, Springer

Examples

#TBA

Weib.reg.BB1.0

Weibull regression under the BB1 copula with known "delta"

Description

See Section 2.6.3 of Emura et al. (2019).

Usage

```
Weib.reg.BB1.0(x.obs,y.obs,dx,dy,zx,zy,delta=0,convergence.par=FALSE)
```

Arguments

x.obs	censored times for X
y.obs	censored times for Y
dx	censoring indicators for X
dy	censoring indicators for Y
zx	matrix of covariates for X
zy	matrix of covariates for Y
delta	known copula parameter ($d \geq 0$)
convergence.par	if TRUE, show the details

Details

Details are seen from the references.

Value

beta_x	regression coefficients for X
beta_y	regression coefficients for Y

Author(s)

Takeshi Emura

References

Emura T, Matsui S, Rondeau V (2019), Survival Analysis with Correlated Endpoints, Joint Frailty-Copula Models, JSS Research Series in Statistics, Springer

Examples

#TBA

Weib.reg.Clayton

Weibull regression under the Clayton copula

Description

See Section 2.6.3 of Emura et al. (2019).

Usage

```
Weib.reg.Clayton(x.obs,y.obs,dx,dy,zx,zy,convergence.par=FALSE)
```

Arguments

x.obs	censored times for X
y.obs	censored times for Y
dx	censoring indicators for X
dy	censoring indicators for Y
zx	matrix of covariates for X
zy	matrix of covariates for Y
convergence.par	if TRUE, show the details

Details

Details are seen from the references.

Value

beta_x	regression coefficients for X
beta_y	regression coefficients for Y

Author(s)

Takeshi Emura

References

Emura T, Matsui S, Rondeau V (2019), Survival Analysis with Correlated Endpoints, Joint Frailty-Copula Models, JSS Research Series in Statistics, Springer

Examples

#TBA

Index

- * **BB1 copula**
 - simu.BB1, 2
 - Weib.reg.BB1, 18
 - Weib.reg.BB1.0, 19
- * **Celebioglu-Cuadras (CC) copula**
 - simu.CC, 4
- * **Clayton copula**
 - simu.Clayton, 5
 - simu.Gumbel, 10
 - Test.Clayton, 11
 - U1.Clayton, 13
 - U2.Clayton, 16
 - Weib.reg.Clayton, 20
- * **FGM copula**
 - simu.FGM, 6
- * **Frank copula**
 - simu.Frank, 7
- * **Goodness-of-fit test**
 - Test.Clayton, 11
 - Test.Gumbel, 12
 - U1.Clayton, 13
 - U1.Gumbel, 15
 - U2.Clayton, 16
 - U2.Gumbel, 17
- * **Gumbel copula**
 - Test.Gumbel, 12
 - U1.Gumbel, 15
 - U2.Gumbel, 17
- * **Gumbel-Barnett (GB) copula**
 - simu.GB, 8
- * **Regression**
 - Weib.reg.BB1, 18
 - Weib.reg.BB1.0, 19
 - Weib.reg.Clayton, 20
- * **Simulation**
 - simu.BB1, 2
 - simu.CC, 4
 - simu.Clayton, 5
 - simu.FGM, 6
- simu.Frank, 7
- simu.GB, 8
- simu.Gumbel, 10
- * **Weibull distribution**
 - simu.BB1, 2
 - simu.CC, 4
 - simu.Clayton, 5
 - simu.FGM, 6
 - simu.Frank, 7
 - simu.GB, 8
 - simu.Gumbel, 10
- * **package**
 - Copula.surv-package, 2
- Copula.surv (Copula.surv-package), 2
- Copula.surv-package, 2
- simu.BB1, 2
- simu.CC, 4
- simu.Clayton, 5
- simu.FGM, 6
- simu.Frank, 7
- simu.GB, 8
- simu.Gumbel, 10
- Test.Clayton, 11
- Test.Gumbel, 12
- U1.Clayton, 13
- U1.Gumbel, 15
- U2.Clayton, 16
- U2.Gumbel, 17
- Weib.reg.BB1, 18
- Weib.reg.BB1.0, 19
- Weib.reg.Clayton, 20